

UDC 547.1-304.2:546.224

R.E. Khoma^{a, *}, *V.M. Baumer*^b, *V.O. Gelmboldt*^c, *N.F. Fedko*^a, *Yu.V. Ishkov*^a,
S.V. Vodzinskii^a, *I.M. Rakipov*^d, *D.V. Menchuk*^a

AMMONIUM SALTS OF SULFUR-CONTAINING OXYANIONS RESULTING FROM REACTION OF SULFUR DIOXIDE WITH AQUEOUS SOLUTIONS OF DIETHYLENTRIAMINE AND BIS(HEXAMETHYLENE)TRIAMINE

^a Odesa I.I. Mechnikov National University, Odesa, Ukraine

^b Institute for Single Crystals, National Academy of Sciences of Ukraine, Kharkiv, Ukraine

^c Odessa National Medical University, Odesa, Ukraine

^d A.V. Bogatsky Physico-Chemical Institute, National Academy of Sciences of Ukraine, Odesa, Ukraine

Interaction in the system «sulfur dioxide–diethylenetriamine (or bis-(hexamethylene)triamine)–water–air oxygen» results in a mixture of double salts (diethylenetriammonium bis(sulfito)-dithionate (Ia) and bis(sulfito)-sulfate trihydrate diethylenetriammonium (Ib)) or bis(hexamethylene)tri ammonium sulfate (II). The obtained ammonium salts of sulfur-containing oxyanions Ia, Ib, and II were characterized by the methods of elemental analysis, X-ray structural analysis, X-ray powder diffraction, IR spectroscopy, and mass spectrometry. Compound Ia crystallizes in monoclinic syngony (space group C2/c, $a=28.1583(5)$ Å, $b=6.84450(10)$ Å, $c=10.8351(2)$ Å, $\beta=93.776(2)^\circ$, $V=2083.71(6)$ Å³, $Z=4$). Compound Ib crystallizes in rhombic syngony (space group Pbcn, $a=18.9777(13)$ Å, $b=10.3131(8)$ Å, $c=11.5414(10)$ Å, $V=2258.9(3)$ Å³, $Z=4$). Compound II crystallizes in triclinic syngony (space group P-1, $a=11.347(3)$ Å, $b=11.700(5)$ Å, $c=18.913(4)$ Å, $\alpha=95.22(3)^\circ$, $\beta=92.52(2)^\circ$, $\gamma=118.27(4)^\circ$, $V=2191.36$ Å³, $Z=2$). The IR spectrum of the mixture of salts Ia and Ib shows the valence vibrations of $\nu(\text{SO})$ of sulfite anion (ν_1 and ν_3), represented by bands at 955, 931, 907, 1026, and 1006 cm⁻¹. Two bands at 566 and 493 cm⁻¹ are the result of the splitting of a doubly degenerate out-of-plane deformation vibration of the ν_4 (E) of dithionate anion. The bands at 1127 and 1078 cm⁻¹ correspond to the oscillations of ν_{as} and ν_{s} of sulfate anion (SO₄²⁻). A decrease in the symmetry of the SO₄²⁻ anion as a result of salt formation is accompanied by the appearance of the band ν_1 (A₁) at 980 cm⁻¹ in its spectrum.

Keywords: sulfur dioxide, 2-hydroxypropylamine, bis(hexamethylene)tri amine, sulfoxidation, spectral characteristics.

DOI: 10.32434/0321-4095-2024-157-6-61-69

Introduction

As is known, sulfur dioxide is one of the main atmospheric pollutants of anthropogenic origin (more than 200 million tons/year): the large-scale emission of this toxicant is accompanied by various negative effects on environmental objects, flora and fauna [1]. Therefore, efforts directed at the development of means for cleaning process gases, sanitary air purification from sulfur dioxide, especially with the use of chemisorption processes by aqueous solutions of organic

chemisorbents based on amines and their derivatives are urgent. At the same time, polyamines, for example, triamines, can be considered as convenient models of polymeric (oligomeric) chemisorbents of the polyethylene polyamine type, on which it is possible to trace the features of chemical transformations in the specified chemisorption systems, namely, salt formation and sulfoxidation reactions. We chose diethylenetriamine (DETA) and bis(hexamethylene)tri amine (BHMT) as model

© R.E. Khoma, V.M. Baumer, V.O. Gelmboldt, N.F. Fedko, Yu.V. Ishkov, S.V. Vodzinskii, I.M. Rakipov, D.V. Menchuk, 2024



This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Ammonium salts of sulfur-containing oxyanions resulting from reaction of sulfur dioxide with aqueous solutions of diethylenetriamine and bis(hexamethylene)tri amine

triamines in this work.

DETA forms ammonium salts (mono-, di-, and triprotonated DETA cations) of the corresponding anions with mineral acids (HCl, HBr, HI, HNO₃, H₂SO₄, H₃PO₄, etc. [2]). Under the conditions of isothermal evaporation of the N(CH₂CH₂NH₂)₂-H₂SO₄-H₂O system, a mixed sulfate-hydrosulfate salt [DETAH₃]₂SO₄(HSO₄) is formed, similar to the AlkNH₂-SO₂-H₂O-O₂ systems (Alk=n-C₇H₁₅ and n-C₈H₁₇)-(AlkNH₃)₃SO₄(HSO₄) [3]. As a result of the interaction of monoprotonated DETA monobromide salt ([DETAH]Br) with sulfur dioxide, a mixed sulfite-hydrobromide salt of triprotonated DETA is formed: [DETAH₃](SO₃)Br [5]. The reaction of DETA with SO₂ in tetradecyl(trihexyl)phosphonium chloride in the presence of air oxygen is accompanied by the formation of a mixed sulfite-chloride salt [DETAH₃](SO₃)Cl, while in the presence of N-methylpyrrolidone and 95% ethanol, sulfite monohydrate [DETAH₂](SO₃)·H₂O is formed with protonated terminal N atoms [6,7].

From an aqueous solution of BHMT with orthophosphoric acid, triammonium hydrogen phosphate BHMT crystallizes in the form of undecahydrate [2]. There is a known case of the monoammonium cation

NH₂(CH₂)₆⁺NH₂(CH₂)₆NH₂ formation in the composition of a polynuclear platinum complex [8].

This work describes the method of synthesis and identification of the reaction products of sulfur dioxide with aqueous solutions of diethylenetriamine and bis(hexamethylene)triamine in the presence of air oxygen, new triammonium salts of diethylenetriammonium bis(sulfite)dithionate (Ia), bis trihydrate diethylenetriammonium (sulfite)-sulfate (Ib) and bis(hexamethylene)triammonium sulfate (II), respectively.

Experimental

A mixture of diethylenetriammonium bis(sulfite)-dithionate (Ia) and diethylenetriammonium bis(sulfite)-sulfate trihydrate (Ib)

A solution of DETA (0.05 mol) in 15 ml of water was poured into a thermostatically controlled cell (20°C) and SO₂ was passed through it in the bubbling mode at a rate of 50 ml·min⁻¹ for 45 min (20°C). The resulting solution was kept at room temperature in air until the water was completely removed. 13.24 g of a mixture of white crystalline products Ia and Ib was isolated.

Calculated, %: C 18.72; H 6.59; N 15.57; S 22.84. Found, %: C 18.11; H 6.30; N 15.84; S 23.03.

MS (EI) I: [DETAH]⁺ (m/z 104, I, 10%);

91, I, 16%; 89, I, 12%; [DETA-NH₃-H]⁺ (85, I, 14%); [DETA-NH₃-2H]⁺ (84, I, 16%); [NH₂CH₂CH₂NHCH₂]⁺ (m/z 73, I, 44%); [SO₂]⁺ (m/z 64, I, 98%); m/z 56, I, 92%; m/z 56, I, 33%; [SO]⁺ (m/z 48, I, 46%); [CH₃CH=NH₂]⁺ (m/z 44, I, 100%); m/z 43, I, 16%; [C₂H₄N]⁺ (m/z 42, I, 31%); [CH₂=NH₂]⁺ (m/z 30, I, 83%).

IR I [ν, cm⁻¹]: 3515 m., 3444 m [ν_{as}(⁺NH)]; 3034 s, 3022 s br, 3012 sh [ν_s(⁺NH)]; 2992 s br, 2984 s br [ν_{as}(CH₂)]; 2938 s br, 2905 s br, 2867 sh [ν_s(CH₂)]; 2735 s br [ν_s(⁺NH), ν_{as}(CH₂)]; 2590 m., 2209 m, 2168 sh, 2135 sh, 2123 m br, 2110 sh, 2076 sh [overtone/combination]; 1633 s [δ_{as}(⁺NH₃)]; 1603s [δ_s(⁺NH₃)]; 1580 sh [δ_{as}(⁺NH₃), δ_{as}(⁺NH₂)]; 1556 sh [δ_s(⁺NH₃), δ_s(⁺NH₂)]; 1525 s [δ_{as}(CH₂)]; 1481 s [δ_s(CH₂)]; 1456 s [ν(C⁺N), δ(C⁺NH)]; 1336 w. [ω(CH₂), ω(⁺NH₃), ω(⁺NH₂)]; 1255 s, 1225 s, 1212 s [ν_{as}(S₂O₆²⁻)]; 1127 s br [ν_{as}(SO₄²⁻), ν_{as}(C⁺NC)]; 1078 s [ν_s(SO₄²⁻), ν(C⁺N)]; 1040 s; 1026 sh, 1006 s [ν_{as,s}(SO₃²⁻)]; 980 s [ν_s(SO₄²⁻), ν_s(SO₃²⁻)]; 955 s [ν_{as}(SO₃²⁻), ν_s(SO₃²⁻), ν_s(CC), δ(⁺NH₃)]; 931 s br [ν_{as}(SO₃²⁻), ν_s(SO₃²⁻), δ(C⁺N)]; 907 s [ν_{as}(SO₃²⁻), ν_s(SO₃²⁻)]; 894 sh; 870 s [ν_s(C-C-N)]; 858 sh; 848 s br [δ(C-C)]; 807 s [ν_s(C-N-C)]; 633 s [δ_{as}(SO₄²⁻)]; 617 s [δ_{as}(SO₄²⁻), δ_s(SO₃²⁻)]; 566 m [δ(S₂O₆²⁻)]; 545 w, 520 sh; 513 s [δ_s(SO₄²⁻)]; 493 s [δ_d(SO₃²⁻)]; 471 w [δ_s(SO₄²⁻), δ(⁺N-C-C-⁺N)]; 458 w [δ(C-⁺N-C)].

Bis(hexamethylene) triammonium sulfate dihydrate (II)

A solution of BHMT (0.01 mol) in 30 ml of water was poured into a thermostatically controlled cell (20°C) and, in the bubbling mode, SO₂ was passed through it at a rate of 50 ml·min⁻¹ for 45 min (20°C). The resulting solution was kept at room temperature in the air until the water was completely removed. The isolated white crystalline product II (4.03 g, yield 97.19%) was used without additional purification. The individuality of the obtained product was confirmed by various physical and chemical methods.

Calculated, %: C 37.76; H 9.13; N 10.93; S 12.76. $C_{24}H_{68}N_6O_{14}S_3$. Found, %: C 37.88; H 9.01; N 11.04; S 12.64. M 761.04.

MS (EI) II: [BHMT-H]⁺ (m/z 214, I, 10%); [BHMD-NH₃]⁺ (m/z 198, I, 12%); m/z 129, I, 36%; m/z 112, I, 41%; [CH₃(CH₂)₄CH=NH₂+H]⁺ (m/z 100, I, 12%); [CH₃(CH₂)₄CH=NH₂]⁺ (m/z 99, I, 10%); [CH₃(CH₂)₄CH=NH₂-H]⁺ (m/z 98, I, 73%); m/z 70, I, 10%; [CH₂=CHNH=CH₂]⁺ (m/z=56, I=19%); m/z 55, I, 18%; [CH₃CH=NH₂]⁺ (m/z 44, I, 21%); m/z 43, I, 10%; [C₂H₄N]⁺ (m/z 42, I, 10%); m/z 41, I, 15%; [CH₂=NH₂]⁺ (m/z 30, I, 100%).

MS (FAB) II: [BHMT+H]⁺ (m/z 216, I, 11%).

IR II [ν, cm⁻¹]: 3436 s [ν(OH)]; 3246 m, 3218 sh, 3000 s, 2963 s, 2944 s, 2928 sh, 2866 m, 2809 m, 2797 m [ν_{as,s}(⁺NH₂), ν_{as,s}(⁺NH₂)]; 2734 w [ν_s(⁺NH), ν_{as}(CH₂)]; 2621 w, 2542 w, 2473 w, 2413 w, 2021 w [overtone/combination]; 1631 m, 1594 w, 1567 sh [δ_{as,s}(⁺NH₃), δ_{as,s}(⁺NH₂)]; 1483 m [δ_s(CH₂)]; 1473 w, 1462 w, 1437 w, 1410 w, 1385 w, 1317 w [ν(C⁺N), δ(C⁺NH)]; 1243 sh, 1221 w [ν_{as}(C-N⁺)]; 1123 s [ν_{as}(SO₄²⁻)]; 1072 sh [ν_s(SO₄²⁻)]; 1061 w, 1049 w, 1030 w [ν_s(C-N⁺)]; 997 w, 973 w, 951 w, 945 w, 908 w. [ρ(⁺NH₃), ρ(⁺NH₂), ν_s(SO₄²⁻), ρ(CH₂)]; 619 s, 583 w [δ_{as}(SO₄²⁻)]; 478 w [δ(⁺NCC), δ_s(SO₄²⁻)].

Analysis for carbon, hydrogen and nitrogen content was carried out on a CHN analyzer; sulfur was determined by the Shoniger technique [9]. IR absorption spectra were recorded on a Perkin-Elmer Spectrum BX II FT-IR System spectrophotometer in 4000–350 cm⁻¹ region as KBr pellets. EI mass spectra were recorded on the MX-1321 device (direct introduction of the sample into the source, energy of ionizing electrons 70 eV). X-ray structural study of the compounds was performed on an Oxford Diffraction Ltd.Xcalibur-3 diffractometer (MoK_α radiation, graphite monochromator, Sapphire-3 CCD detector). X-ray powder diffraction (XRD) was accomplished on a Siemens D500 powder diffractometer (Bragg-Brentano geometry, CuK_α radiation, Ni filter).

Results and discussion

Mass spectra of the mixture of salts Ia and Ib and of compound II show defragmentation typical for

1-n-alkylamines [10] with the formation of the ion [CH₂=NH₂]⁺, the peak of which has the maximum intensity for compound II. The characteristics of the fragmentation products of DETA in the mass spectrum of the mixture of its ammonium salts Ia and Ib and the tabulated mass spectrum of DETA [10] are in good agreement.

The main crystallographic data and refinement results are given in Table 1. Bond lengths and valence angles in the structures are given in Tables 2 and 3. It can be seen that these characteristics have the usual values for similar compounds (for example ref. [4]). In structure Ia, the dithionate ion is in a private position in the center of symmetry (Fig. 1). In structure Ib, the S(2)–O(4) bond of the sulfate ion is on the second-order axis (Fig. 2), so the remaining oxygen atoms of this ion are disordered. In addition, in this structure, the water molecule O(9) is also on the second-order axis. The configuration of the cation in structures Ia and Ib is markedly different. Thus, in Ib, the non-hydrogen atoms of the cation backbone lie almost in one plane (standard deviation is 0.094 Å), and in Ia the terminal nitrogen atoms deviate from the mean plane (standard deviation is 0.056 Å) of the other core atoms (N(2) and N(3) at 1.248 Å and 1.239 Å, respectively). These conformational differences can also be illustrated by data on torsion angles (Table 4): they differ little from +180° or –180° for the almost flat structure Ib, while these differences are significant in Ia. Layered packing is observed in both studied structures (Figs. 3 and 4). In the Ia structure, anions are arranged in layers in (001) planes, cations are located between layers of anions and form layers as well. A similar packing is observed in salt Ib: anions together with water crystallization molecules form layers parallel to (010), and layers of cations are located between them. With such a packing, a three-dimensional system of hydrogen bonds is observed in both structures, data on which are given in Tables 5 and 6. Some of these bonds are «bifurcated», or three-centered: N(2)–H(2E)···O(5) and N(2)–H(2E)···O(5)^{-x+1,-y,-z+1} in Ia, N(1)–H(1A)···O(2)^{x,-y,z+1/2}, N(1)–H(1A)···O(1)^{x,-y,z+1/2}, N(2)–H(2A)···O(2)^{-x+1/2,y+1/2,z}, N(2)–H(2A)···O(3)^{-x+1/2,y+1/2,z}, N(2)–H(2C)···O(5)^{-x+1/2,-y+1/2,z-1/2}, N(2)–H(2C)···O(6)^{x-1/2,-y+1/2,-z+1}, N(3)–H(3A)···O(6) and N(3)–H(3A)···O(7)^{-x+1,y,-z+3/2} in Ib. It should be noted that they are not formed in equal amounts in a mixture of compounds Ia and Ib. According to the calculation of the powder X-ray pattern by the Rietveld method using as initial data the structural models obtained by single-crystal method (Fig. 5), the reaction product contains 81.4 wt.% salt (Ia) and 18.6% salt (Ib). No other phases were found in the powder sample.

Table 1

Crystallographic data, X-ray diffraction conditions, and structure refinement characteristics for Ia and Ib

Characteristic	Values and types	
	Ia	Ib
molecular formula	C ₈ H ₃₂ N ₆ O ₁₂ S ₄	C ₈ H ₃₈ N ₆ O ₁₃ S ₃
M _r	532.64	522.62
T, K	293(2)	293(2)
syngony	monoclinic	rhombic
space class	C2/c	Pbcn
a, Å	28.1583(5)	18.9777(13)
b, Å	6.84450(10)	10.3131(8)
c, Å	10.8351(2)	11.5414(10)
β, deg	93.776(2)	–
V, Å ³	2083.71(6)	2258.9(3)
Z	4	4
ρ, g/cm ³	1.698	1.537
μ(MoK _α), mm ⁻¹	0.527	0.399
θ range, deg	3.69–31.78	3.41–32.32
crystal dimensions, mm	0.35×0.20×0.15	0.40×0.25×0.15
F ₀₀₀	1128	1120
T _{min} /T _{max}	0.837/0.925	0.8568/0.9426
number of reflections: measured	11832	22184
independent with 3 I _{hkl} >2σ(I)	3149	3645
R _{int}	0.0560	0.0488
fullness, %	95.1	96.9
number of refined parameters	141	169
R _F /wR ² on observed reflections	0.0451/0.1241	0.0391/0.1077
R _F /wR ² on independent reflections	0.0495/0.1304	0.0490/0.1156
S	0.992	0.982
Δρ _{min} /Δρ _{max} , e/Å ³	–0.666/0.424	–0.386/0.472

Table 2

Bond lengths and bond angles in the structure Ia*

Bond	d, Å	Bond	d, Å	Bond	d, Å
S(1)–O(2)	1.5194(12)	S(2)–O(5)	1.4799(13)	N(2)–C(2)	1.491(2)
S(1)–O(1)	1.5223(12)	S(2)–S(2)#1	2.1853(9)	N(3)–C(4)	1.481(2)
S(1)–O(3)	1.5306(12)	N(1)–C(1)	1.4887(16)	C(1)–C(2)	1.511(2)
S(2)–O(6)	1.367(2)	N(1)–C(3)	1.4951(17)	C(3)–C(4)	1.5092(19)
S(2)–O(4)	1.4709(14)				
Angle	ω, deg	angle	ω, deg		
O(2)–S(1)–O(1)	105.47(7)	O(4)–S(2)–S(2)#1	102.73(6)		
O(2)–S(1)–O(3)	103.94(7)	O(5)–S(2)–S(2)#1	101.76(7)		
O(1)–S(1)–O(3)	104.47(7)	C(1)–N(1)–C(3)	111.60(10)		
O(6)–S(2)–O(4)	126.35(14)	N(1)–C(1)–C(2)	111.98(11)		
O(6)–S(2)–O(5)	108.37(13)	N(2)–C(2)–C(1)	112.71(12)		
O(4)–S(2)–O(5)	112.51(8)	N(1)–C(3)–C(4)	111.62(11)		
O(6)–S(2)–S(2)#1	101.32(11)	N(3)–C(4)–C(3)	112.91(12)		

Note: * – symmetric transformation to obtain equivalent atoms #1 are as follows: $-x+1$, $-y+1$, $-z+1$.

Table 3

Bond lengths and bond angles in the structure Ib*

Bond	d, Å	Bond	d, Å	Bond	d, Å
S(1)–O(1)	1.5093(9)	S(2)–O(7)#1	1.427(2)	N(1)–C(1)	1.4785(14)
S(1)–O(3)	1.5112(9)	S(2)–O(7)	1.427(2)	N(2)–C(2)	1.4716(14)
S(1)–O(2)	1.5331(9)	S(2)–O(4)	1.4555(12)	N(3)–C(4)	1.4599(16)
S(2)–O(5)	1.4256(17)	S(2)–O(6)	1.495(2)	C(1)–C(2)	1.5040(16)
S(2)–O(5)#1	1.4257(17)	S(2)–O(6)#1	1.495(2)	C(3)–C(4)	1.5053(16)
S(2)–O(7)#1	1.427(2)	N(1)–C(3)	1.4757(14)		
angle	ω , deg	angle	ω , deg		
O(1)–S(1)–O(3)	105.82(5)	O(4)–S(2)–O(6)	107.03(9)		
O(1)–S(1)–O(2)	104.60(5)	O(5)#1–S(2)–O(6)#1	107.70(14)		
O(3)–S(1)–O(2)	104.44(5)	O(7)#1–S(2)–O(6)#1	105.42(16)		
O(5)#1–S(2)–O(7)#1	114.80(14)	O(7)#1–S(2)–O(6)#1	105.42(16)		
O(5)–S(2)–O(7)	114.80(14)	C(3)–N(1)–C(1)	111.55(9)		
O(5)–S(2)–O(4)	111.65(9)	N(1)–C(1)–C(2)	110.62(9)		
O(7)–S(2)–O(4)	109.74(11)	N(2)–C(2)–C(1)	109.39(9)		
O(5)–S(2)–O(6)	107.70(14)	N(1)–C(3)–C(4)	109.70(10)		
O(7)–S(2)–O(6)	105.43(16)	N(3)–C(4)–C(3)	110.10(10)		

Note: * – symmetric transformation to obtain equivalent atoms #1 are as follows: $-x+1, y, -z+3/2$.

Table 4
Torsion angles in the cation for Ia and Ib

Bond	Torsion angle	
	Ia	Ib
C(3)–N(1)–C(1)–C(2)	-173.60(11)	-178.66(10)
N(1)–C(1)–C(2)–N(2)	-68.78(15)	-178.34(9)
C(1)–N(1)–C(3)–C(4)	173.81(11)	-172.60(11)
N(1)–C(3)–C(4)–N(3)	71.13(14)	169.72(10)




Fig. 1. Atom numbering scheme and thermal ellipsoids (probability level 50%) for the structure $[(\text{NH}_3\text{CH}_2\text{CH}_2)_2\text{NH}]_2(\text{SO}_3)_2\text{S}_2\text{O}_6$




Fig. 2. Atom numbering scheme and thermal ellipsoids (probability level 50%) for the structure $[(\text{NH}_3\text{CH}_2\text{CH}_2)_2\text{NH}]_2(\text{SO}_3)_2\text{SO}_4 \cdot 3\text{H}_2\text{O}$




Fig. 3. Projection xy_0 of the structure $[(\text{NH}_3\text{CH}_2\text{CH}_2)_2\text{NH}]_2(\text{SO}_3)_2\text{S}_2\text{O}_6$. Hydrogen bonds are represented by dashed lines

Table 5

Geometrical characteristics of hydrogen bonds in the structure Ia

Contact D–H...A	Distance, Å			DHA angle, deg	Transformation for atom A
	d(D–H)	d(H...A)	d(D...A)		
N(1)–H(1A)···O(2)#2	0.90	1.94	2.8001(15)	160.3	x, y–1, z
N(1)–H(1B)···O(1)#3	0.90	1.85	2.7448(16)	177.1	x, –y+1, z+1/2
N(2)–H(2C)···O(4)#2	0.83	2.17	2.9672(18)	160.0	x, y–1, z
N(2)–H(2D)···O(3)#2	0.83	1.99	2.7291(18)	147.1	x, y–1, z
N(2)–H(2E)···O(5)	0.83	2.03	2.7964(18)	154.3	
N(2)–H(2E)···O(5)#4	0.83	2.60	3.0951(19)	119.3	–x+1, –y, –z+1
N(3)–H(3C)···O(1)#5	0.88	1.95	2.7863(17)	157.3	–x+3/2, –y+3/2, –z+1
N(3)–H(3D)···O(3)	0.88	1.88	2.7325(17)	163.2	
N(3)–H(3E)···O(2)#2	0.88	1.93	2.7740(17)	160.3	x, y–1, z

Table 6

Geometrical characteristics of hydrogen bonds in the structure Ib

Contact D–H...A	Distance, Å			DHA angle, deg	Transformation for atom A
	d(D–H)	d(H...A)	d(D...A)		
O(8)–H(8A)···O(5)	0.840(8)	2.014(8)	2.850(3)	173.5(13)	
O(8)–H(8A)···O(7)#1	0.840(8)	2.171(9)	2.918(3)	148.2(12)	–x+1, y, –z+3/2
O(8)–H(8B)···O(2)#2	0.839(7)	2.046(7)	2.8374(16)	157.2(15)	–x+1/2, –y+1/2, z+1/2
O(9)–H(9)···O(3)	0.841(8)	2.035(8)	2.8701(13)	171.5(16)	
N(1)–H(1A)···O(2)#3	0.89	1.87	2.7384(12)	167.1	x, –y, z+1/2
N(1)–H(1A)···O(1)#3	0.89	2.58	3.1813(13)	125.3	x, –y, z+1/2
N(1)–H(1B)···O(3)	0.89	1.82	2.6948(12)	170.3	
N(2)–H(2A)···O(2)#4	0.85	1.99	2.8405(13)	174.3	–x+1/2, y+1/2, z
N(2)–H(2A)···O(3)#4	0.85	2.59	3.1108(13)	120.2	–x+1/2, y+1/2, z
N(2)–H(2B)···O(1)#2	0.85	1.94	2.7675(13)	164.9	–x+1/2, –y+1/2, z+1/2
N(2)–H(2C)···O(5)#5	0.85	1.85	2.642(2)	153.9	–x+1/2, –y+1/2, z–1/2
N(2)–H(2C)···O(6)#6	0.85	2.16	2.913(3)	146.7	X–1/2, –y+1/2, –z+1
N(3)–H(3A)···O(6)	0.92	1.85	2.717(3)	156.1	
N(3)–H(3A)···O(7)#1	0.92	1.96	2.822(3)	154.5	–x+1, y, –z+3/2
N(3)–H(3B)···O(4)#7	0.92	1.98	2.8399(13)	154.6	–x+1, –y+1, –z+1
N(3)–H(3C)···O(1)#8	0.92	1.91	2.8184(14)	170.7	–x+1, y, –z+1/2

Indication of the diffractogram lines for compound II according to the Rietveld method in the line assignment mode showed that this substance belongs to triclinic class (space group P-1, $a=11.347(3)$ Å, $b=11.00(5)$ Å, $c=18.913(4)$ Å, $\alpha=95.22(3)^\circ$, $\beta=92.52(2)^\circ$, $\gamma=118.27(4)^\circ$, $V=2191.36$ Å³, $Z=2$).

The assignment of bands in IR spectra I and II for cations [DETAH₃]³⁺ and [BHMTH₃]³⁺ was made taking into account the literature data [3,11,12]. The bands observed in the IR spectrum of I in the region of 3515–2867 cm⁻¹ refer to symmetric and asymmetric valence vibrations $\nu(\overset{+}{\text{N}}\text{H}_3)$, $\nu(\overset{+}{\text{N}}\text{H}_2)$ and $\nu(\text{CH}_2)$. The bands located between 1633 and 1456 cm⁻¹ correspond

to the scissoring vibrations of the $\overset{+}{\text{N}}\text{H}_3$, $\overset{+}{\text{N}}\text{H}_2$ and CH₂ groups, and to the valence $\nu(\text{C–N})$ and deformation $\delta(\text{CNH})$ vibrations.

A broad intense band in the IR spectrum of dihydrate II with seven well-defined maxima in the region of 3440–2730 cm⁻¹ is associated with vibrations of $\nu(\text{OH})$ and $\nu(\text{NH})$ molecules of water and triammonium cation [BHMTH₃]³⁺. In the interval of 1750–1500 cm⁻¹, a doublet is observed at 1631 and 1694 cm⁻¹ with a shoulder at 1567 cm⁻¹, which probably refers to overlapping scissor vibrations of NH groups and water molecules characteristic for this region, similarly [HEEDA]. A multiplet of medium and weak intensity in the region of 1483–1317 cm⁻¹




Fig. 4. Projection xy_0 of the structure $[(\text{NH}_3\text{CH}_2\text{CH}_2)_2\text{NH}_2]_2(\text{SO}_3)_2\text{SO}_4 \cdot 3\text{H}_2\text{O}$. Hydrogen bonds are shown by dashed lines




Fig. 5. Rietveld refinement for powder diffractogram of sample I. Dots show experimental curve, solid line displays calculation. Arrays of vertical bars show diffraction maxima (upper row for Ia, and lower row for Ib). The lower curve displays the difference between experimental and calculated intensity values in each point



can be attributed to mixed valence-deformation vibrations $\nu(\text{C}_\text{N}^+)$ and $\delta(\text{CH})$.

In the IR spectrum of the mixture of salts Ia and Ib, the valence vibrations $\nu(\text{SO})$ of the SO_3^{2-} (ν_1 and ν_3 , respectively) are represented by intense bands at 955 cm^{-1} , 931 cm^{-1} and 907 cm^{-1} ; there are also bands of lower intensity at 1026 and 1006 cm^{-1} in this region. Two bands of medium and strong intensity

at 566 and 493 cm^{-1} can be considered as the result of the splitting of the doubly degenerate out-of-plane deformation vibration ν_4 (E) of $\text{S}_2\text{O}_6^{2-}$ ion. There are strong bands at 1127 and 1078 cm^{-1} in the IR spectrum of I, corresponding to vibrations ν_{as} and $\nu_s(\text{SO}_4^{2-})$, respectively. The increased intensity of these bands (the strongest in the spectrum) is obviously due to the fact that vibrations $\nu(\text{C}_\text{N}^+)$ also contribute to them. The decrease in symmetry of anion as a result of salt formation is accompanied by the appearance of a strong band of fully symmetric valence vibration ν_1 (A_1) at 980 cm^{-1} . The salt with BHMT has also the most intense band at 1123 cm^{-1} ($\nu_{\text{as}}(\text{SO}_4^{2-})$); and $\nu_1(\text{A}_1)$ is observed as a band of medium intensity at 980 cm^{-1} .

The IR spectrum of the mixture of Ia and Ib, similarly to results presented in ref. [4], has the bands in the region around 1210 – 1255 cm^{-1} and 570 cm^{-1} , which can be attributed to $\nu_{\text{as}}(\text{S}_2\text{O}_6^{2-})$ and $\delta(\text{S}_2\text{O}_6^{2-})$, respectively, which confirms the presence in the obtained products along with bis-sulfite (Ia) and (bis)sulfite-sulfate (Ib) anions of dithionate anions.

Thus, taking into account the data of [13], chemisorption of SO_2 by an aqueous solution of DETA produces ammonium hydrosulfite $[\text{H}_3\text{NCH}_2\text{CH}_2\text{NHCH}_2\text{CH}_2\text{NH}_3](\text{HSO}_3)_2$, in which terminal nitrogen atoms are obviously protonated, similar to sulfite $\text{H}_3\text{NCH}_2\text{CH}_2\text{NHCH}_2\text{CH}_2\text{NH}_3(\text{SO}_3) \cdot \text{H}_2\text{O}$ [6]. In the process of keeping the SO_2 –DETA– H_2O reaction mixture in air, two mixed salts are formed as follows:



and the degree of conversion by reaction (1) is 4.3 times greater than by reaction (2). In the case of lipophilic BHMT ($\log P_{\text{ow}} = 1.30$), sulfoxidation process $\text{S(IV)} \rightarrow \text{S(VI)}$ runs more completely, like hexamethylenediamine [9] ($\log P_{\text{ow}} = 0.40$), as opposed to hydrophilic DETA ($\log P_{\text{ow}} = -1.30$).

¹ Reaction mass of 7-azatridecane-1,13-diamine and hexamethylenediamine. Available from: <https://echa.europa.eu/bg/registration-dossier/-/registered-dossier/14602/4/8/>.

² Chemical Book. Available from: <https://www.chemicalbook.com/msds/>.

REFERENCES

1. Zhang K., Wang X. Pollution haven hypothesis of global CO₂, SO₂, NO_x—evidence from 43 economies and 56 sectors // *Int. J. Environ. Res. Public Health*. — 2021. — Vol.18. — No. 12. — Art. No. 6552.
2. Ilioudis C.A., Georganopoulou D.G., Steed J.W. Insights into supramolecular design 2: analysis of anion coordination geometry of oxoanions in a protonated polyamine matrix // *CrystEngComm*. — 2002. — Vol.4. — No. 6. — P.26-36.
3. Crystal structure, phase transitions and ir spectroscopic investigations, of a new sulfate templated by diethylenetriammonium / Ettis H., Hajji M., Touati R., Guerfel T. // *Phosphorus Sulfur Silicon Relat. Elem.* — 2016. — Vol.191. — No. 5. — P.765-771.
4. Onium salts of sulfur-containing oxyanions resulting from reaction of sulfur(IV) oxide with aqueous solutions of 1,2-diamines and morpholine / Khoma R.E., Gel'mbol'dt V.O., Ennan A.A., Baumer V.N., Puzan A.N., Koksharova T.V., et al. // *Russ. J. Inorg. Chem.* — 2017. — Vol.62. — No. 6. — P.736-745.
5. The reactivity of diethylenetriamine (DETA) and its simple salts with CO₂ and SO₂ / Doyle K.A., Murphy L.J., Robertson K.N., Clyburne J.A.C. // *Abstracts of the 96th Canadian Chemistry Conference and Exhibition*. — Canada, Quebec, 2013. — P.00871.
6. Ionic liquids and acid gas capture: water and oxygen as confounding factors / Murphy L.J., McPherson A.M., Robertson K.N., Clyburne J.A.C. // *Chem. Commun.* — 2012. — Vol.48. — No. 9. — P.1227-1229.
7. Liquid-solid phase-change absorption of acidic gas by polyamine in nonaqueous organic solvent. / Zhao W., Zhao Q., Zhang Z., Liu J., Chen R., Chen Y., et al. // *Fuel*. — 2017. — Vol.209. — P.69-75.
8. Solution studies of dinuclear polyamine-linked platinum-based antitumour complexes. / Ruhayel R.A., Zgani I., Berners-Price S.J., Farrell N.P. // *Dalton Trans.* — 2011. — Vol.40. — No. 16. — P.4147-4154.
9. Products of interaction between sulfur(IV) oxide and aqueous solutions of hexamethylenediamine and tert-butylamine: the crystal structure of hexamethylenediammonium sulfate dihydrate / Khoma R.E., Ennan A.A., Shishkin O.V., Baumer V.N., Gel'mbol'dt V.O. // *Russ. J. Inorg. Chem.* — 2012. — Vol.57. — No. 12. — P.1559-1562.
10. Vul'fson N.S., Zaikin V.G., Mikaya A.I. Mass spectrometry of organic compounds. — M.: Khimiya, 1986. — 312 p.
11. Bis(hexamethylene)triamine as potential shale inhibitor in water-based drilling fluid / Zhong H., Qiu Z., Huang W., Xie B., Wang W. // *Open Pet. Eng. J.* — 2013. — Vol.12. — No. 6. — P.49-56.
12. Crystal structure, vibrational spectra and non-linear optical properties of diethylenetriammonium hexabromobismuthate: C₄H₁₆N₃BiBr₆ / Dammak H., Feki H., Boughzala H., Abid Y. // *Spectrochim. Acta A Mol. Biomol. Spectrosc.* — 2015. — Vol.137. — P.1235-1243.
13. Khoma R.E., Dlubovskii R.M., Gelmboldt V.O. Chemisorption of sulfur dioxide by aqueous solutions of ethanolamines under static conditions // *Russ. J. Gen. Chem.* — 2016. — Vol.86. — No. 8. — P.1811-1818.

Received 25.07.2024

АМОНІЄВІ СОЛІ СІРКОВМІСНИХ ОКСИАНІОНІВ — ПРОДУКТИ ВЗАЄМОДІЇ ДІОКСИДУ СІРКИ З ВОДНИМИ РОЗЧИНАМИ ДІЕТИЛЕНТРИАМІНУ ТА БІС(ГЕКСАМЕТИЛЕН)ТРИ АМІНУ

Р.Є. Хома, В.М. Баумер, В.О. Гельмбольдт, Н.Ф. Федько, Ю.В. Ішков, С.В. Водзінський, І.М. Ракіпов, Д.В. Менчук

При взаємодії в системі «діоксид сірки—діетилентриамін (або біс-(гексаметилен)триамін)—вода—кисень повітря» утворюються суміші подвійних солей (біс(сульфіто)-дитіонату діетилентриамонію (Ia) та тригідрату біс(сульфіто)-сульфату діетилентриамонію (Ib)) або сульфат біс(гексаметилен)триамонію (II). Одержані амонієві солі сірковмісних оксианіонів Ia, Ib та II охарактеризовано методами елементного аналізу, РСА, РФА, ІЧ-спектроскопії та мас-спектрометрії. Сполука Ia кристалізується у моноклінній сингонії (просторова група C2/c, a=28.1583(5) Å, b=6.84450(10) Å, c=10.8351(2) Å, β=93.776(2)°, V=2083.71(6) Å³, Z=4), Ib — у ромбічній (просторова група Rbcn, a=18.9777(13) Å, b=10.3131(8) Å, c=11.5414(10) Å, V=2258.9(3) Å³, Z=4), а II — триклінній (просторова група P-1, a=11,347(3) Å, b=11,700(5) Å, c=18,913(4) Å, α=95.22(3)°, β=92,52(2)°, γ=118,27(4)°, V=2191,36 Å³, Z=2). В ІЧ-спектрі суміші солей Ia та Ib валентні коливання ν(SO) сульфат аніона (ν₁ та ν₃) надані смугами при 955, 931, 907, 1026 та 1006 см⁻¹. Дві смуги при 566 та 493 см⁻¹ — це результат розщеплення двічі виродженого позаплощинного деформаційного коливання ν₄ (E) дитіонат аніона. Смуги при 1127 та 1078 см⁻¹ відповідають коливанням ν_{as} та ν_s (SO₄²⁻). Зниження симетрії аніона SO₄²⁻ в результаті солеутворення супроводжується появою в його спектрі смуги ν₁ (A₁) при 980 см⁻¹.

Ключові слова: діоксид сірки; 2-гідроксипропіламін; біс(гексаметилен)триамін; сульфоокиснення; спектральні характеристики.

AMMONIUM SALTS OF SULFUR-CONTAINING OXYANIONS RESULTING FROM REACTION OF SULFUR DIOXIDE WITH AQUEOUS SOLUTIONS OF DIETHYLENTRIAMINE AND BIS(HEXAMETHYLENE)TRIAMINE

R.E. Khoma ^{a,*}, V.M. Baumer ^b, V.O. Gelmboldt ^c,
N.F. Fedko ^a, Yu.V. Ishkov ^a, S.V. Vodzinskii ^a, I.M. Rakipov ^d,
D.V. Menchuk ^a

^a Odesa I.I. Mechnikov National University, Odesa, Ukraine

^b Institute for Single Crystals, National Academy of Sciences of Ukraine, Kharkiv, Ukraine

^c Odessa National Medical University, Odesa, Ukraine

^d A.V. Bogatsky Physico-Chemical Institute, National Academy of Sciences of Ukraine, Odesa, Ukraine

* e-mail: rek@onu.edu.ua

Interaction in the system «sulfur dioxide–diethylenetriamine (or bis-(hexamethylene)triamine)–water–air oxygen» results in a mixture of double salts (diethylenetriammonium bis(sulfite)-dithionate (Ia) and bis(sulfite)-sulfate trihydrate diethylenetriammonium (Ib) or bis(hexamethylene)triammonium sulfate (II). The obtained ammonium salts of sulfur-containing oxyanions Ia, Ib, and II were characterized by the methods of elemental analysis, X-ray structural analysis, X-ray powder diffraction, IR spectroscopy, and mass spectrometry. Compound Ia crystallizes in monoclinic syngony (space group C2/c, $a=28.1583(5)$ Å, $b=6.84450(10)$ Å, $c=10.8351(2)$ Å, $\beta=93.776(2)^\circ$, $V=2083.71(6)$ Å³, $Z=4$). Compound Ib crystallizes in rhombic syngony (space group Pbcn, $a=18.9777(13)$ Å, $b=10.3131(8)$ Å, $c=11.5414(10)$ Å, $V=2258.9(3)$ Å³, $Z=4$). Compound II crystallizes in triclinic syngony (space group P-1, $a=11.347(3)$ Å, $b=11.700(5)$ Å, $c=18.913(4)$ Å, $\alpha=95.22(3)^\circ$, $\beta=92.52(2)^\circ$, $\gamma=118.27(4)^\circ$, $V=2191.36$ Å³, $Z=2$). The IR spectrum of the mixture of salts Ia and Ib shows the valence vibrations of ν (SO) of sulfite anion (ν_1 and ν_3), represented by bands at 955, 931, 907, 1026, and 1006 cm⁻¹. Two bands at 566 and 493 cm⁻¹ are the result of the splitting of a doubly degenerate out-of-plane deformation vibration of the ν_4 (E) of dithionate anion. The bands at 1127 and 1078 cm⁻¹ correspond to the oscillations of ν_{as} and ν_s of sulfate anion (SO₄²⁻). A decrease in the symmetry of the SO₄²⁻ anion as a result of salt formation is accompanied by the appearance of the band ν_1 (A₁) at 980 cm⁻¹ in its spectrum.

Keywords: sulfur dioxide; 2-hydroxypropylamine; bis(hexamethylene)triamine; sulfoxidation; spectral characteristics.

REFERENCES

- Zhang K, Wang X. Pollution haven hypothesis of global CO₂, SO₂, NO_x—evidence from 43 economies and 56 sectors. *Int J Environ Res Public Health*. 2021; 18(12): 6552. doi: 10.3390/ijerph18126552.
- Ilioudis CA, Georganopoulou DG, Steed JW. Insights into supramolecular design 2: analysis of anion coordination geometry of oxoanions in a protonated polyamine matrix. *CrystEngComm*. 2002; 4: 26–36. doi: 10.1039/B109449A.
- Ettis H, Hajji M., Touati R., Guerfel T. Crystal structure, phase transitions and IR spectroscopic investigations of a new sulfate templated by diethylenetriammonium. *Phosphorus Sulfur Silicon Relat Elem*. 2016; 191(5): 765–771. doi: 10.1080/10426507.2015.1100181.
- Khoma RE, Gel'mbol'dt VO, Ennan AA, Baumer VN, Puzan AN, Koksharova TV, et al. Onium salts of sulfur-containing oxyanions resulting from reaction of sulfur(IV) oxide with aqueous solutions of 1,2-diamines and morpholine. *Russ J Inorg Chem*. 2017; 62: 736–745. doi: 10.1134/S0036023617060109.
- Doyle KA, Murphy LJ, Robertson KN, Clyburne JAC. The reactivity of diethylenetriamine (DETA) and its simple salts with CO₂ and SO₂. *Abstracts of the 96th Canadian Chemistry Conference and Exhibition*; May 26–30, 2013; Canada, Quebec. 2013. p. 00871.
- Murphy LJ, McPherson AM, Robertson KN, Clyburne JAC. Ionic liquids and acid gas capture: water and oxygen as confounding factors. *Chem Commun*. 2012; 48(9): 1227–1229. doi: 10.1039/c1cc16008g.
- Zhao W, Zhao Q, Zhang Z, Liu J, Chen R, Chen Y, et al. Liquid-solid phase-change absorption of acidic gas by polyamine in nonaqueous organic solvent. *Fuel*. 2017; 209: 69–75. doi: 10.1016/j.fuel.2017.07.081.
- Ruhayel RA, Zgani I, Berners-Price SJ, Farrell NP. Solution studies of dinuclear polyamine-linked platinum-based antitumour complexes. *Dalton Trans*. 2011; 40: 4147–4154. doi: 10.1039/c1dt00001b.
- Khoma RE, Ennan AA, Shishkin OV, Baumer VN, Gel'mbol'dt VO. Products of interaction between sulfur(IV) oxide and aqueous solutions of hexamethylenediamine and tert-butylamine: the crystal structure of hexamethylenediammonium sulfate dihydrate. *Russ J Inorg Chem*. 2012; 57: 1559–1562. doi: 10.1134/S003602361212008X.
- Vul'fson NS, Zaikin VG, Mikaya AI. *Mass spectrometry of organic compounds*. Moscow: Chemistry; 1986. 312 p. (in Russian).
- Zhong H, Qiu Z, Huang W, Xie B, Wang W. Bis(hexamethylene)triamine as potential shale inhibitor in water-based drilling fluid. *Open Pet Eng J*. 2013; 6: 49–56. doi: 10.2174/1874834101306010049.
- Dammak H, Feki H, Boughzala H, Abid Y. Crystal structure, vibrational spectra and non-linear optical properties of diethylenetriammonium hexabromobismuthate: C₄H₁₆N₃BiBr₆. *Spectrochim Acta A Mol Biomol Spectrosc*. 2015; 137: 1235–1243. doi: 10.1016/j.saa.2014.08.150.
- Khoma RE, Dlubovskii RM, Gelmboldt VO. Chemisorption of sulfur dioxide by aqueous solutions of ethanalamines under static conditions. *Russ J Gen Chem*. 2016; 86: 1811–1818. doi: 10.1134/S1070363216080065.